

RESEARCH

Open Access

Transforming growth factor β 1-induced astrocyte migration is mediated in part by activating 5-lipoxygenase and cysteinyl leukotriene receptor 1

Xue-Qin Huang, Xia-Yan Zhang, Xiao-Rong Wang, Shu-Ying Yu, San-Hua Fang, Yun-Bi Lu, Wei-Ping Zhang and Er-Qing Wei*

Abstract

Background: Transforming growth factor- β 1 (TGF- β 1) is an important regulator of cell migration and plays a role in the scarring response in injured brain. It is also reported that 5-lipoxygenase (5-LOX) and its products, cysteinyl leukotrienes (CysLTs, namely LTC₄, LTD₄ and LTE₄), as well as cysteinyl leukotriene receptor 1 (CysLT₁R) are closely associated with astrocyte proliferation and glial scar formation after brain injury. However, how these molecules act on astrocyte migration, an initial step of the scarring response, is unknown. To clarify this, we determined the roles of 5-LOX and CysLT₁R in TGF- β 1-induced astrocyte migration.

Methods: In primary cultures of rat astrocytes, the effects of TGF- β 1 and CysLT receptor agonists on migration and proliferation were assayed, and the expression of 5-LOX, CysLT receptors and TGF- β 1 was detected. 5-LOX activation was analyzed by measuring its products (CysLTs) and applying its inhibitor. The role of CysLT₁R was investigated by applying CysLT receptor antagonists and CysLT₁R knockdown by small interfering RNA (siRNA). TGF- β 1 release was assayed as well.

Results: TGF- β 1-induced astrocyte migration was potentiated by LTD₄, but attenuated by the 5-LOX inhibitor zileuton and the CysLT₁R antagonist montelukast. The non-selective agonist LTD₄ at 0.1 to 10 nM also induced a mild migration; however, the selective agonist N-methyl-LTC₄ and the selective antagonist Bay cysLT2 for CysLT₂R had no effects. Moreover, CysLT₁R siRNA inhibited TGF- β 1- and LTD₄-induced astrocyte migration by down-regulating the expression of this receptor. However, TGF- β 1 and LTD₄ at various concentrations did not affect astrocyte proliferation 24 h after exposure. On the other hand, TGF- β 1 increased 5-LOX expression and the production of CysLTs, and up-regulated CysLT₁R (not CysLT₂R), while LTD₄ and N-methyl-LTC₄ did not affect TGF- β 1 expression and release.

Conclusions: TGF- β 1-induced astrocyte migration is, at least in part, mediated by enhanced endogenous CysLTs through activating CysLT₁R. These findings indicate that the interaction between the cytokine TGF- β 1 and the pro-inflammatory mediators CysLTs in the regulation of astrocyte function is relevant to glial scar formation.

Keywords: Transforming growth factor- β 1, Cysteinyl leukotriene, Cysteinyl leukotriene receptor, 5-lipoxygenase, Astrocyte, migration, Glial scar

* Correspondence: weieq2006@zju.edu.cn
Department of Pharmacology, Key Laboratory of Medical Neurobiology of Ministry of Health of China, Zhejiang Province Key Laboratory of Neurobiology, Zhejiang University School of Medicine, 866 Yuhangtang Road, Hangzhou 310058, China

Background

Glial scar formation is a critical event in repair responses after injury of the central nervous system (CNS) [1,2]. The glial scar is a complex of cellular components and mainly consists of reactive astrocytes (undergoing proliferation and morphological changes). Following focal CNS injury, reactive astrocytes migrate towards the lesion and then organize into a densely packed glial scar [1,2]. As the key step of glial scar formation, astrocyte migration is regulated by various factors [3-5], among which transforming growth factor- β (TGF- β) is known as an important regulator [5,6].

TGF- β , a family of multifunctional cytokines, regulates a broad diversity of physiological and pathological processes, including wound healing, inflammation, cell proliferation, differentiation, migration and extracellular matrix synthesis [7-10]. TGF- β 1 is an important mediator in the pathogenesis of several disorders in the CNS, such as in the organization of a glial scar in response to injury and in several neurodegenerative disorders [7,11,12]. After CNS injury, elevated TGF- β levels in astrocytes have been shown to induce astrocytic scar formation [13], and are also associated with ischemic brain injury [14,15].

On the other hand, cysteinyl leukotrienes (CysLTs, namely LTC₄, LTD₄, and LTE₄), the 5-lipoxygenase (5-LOX, EC 1.13.11.34) metabolites of arachidonic acid [16], are bioactive lipid mediators that modulate immune and inflammatory responses [16-19] through activating their receptors, CysLT₁R and CysLT₂R [17,20,21]. In the rat brain, 5-LOX is activated and the production of CysLTs is enhanced after focal cerebral ischemia, resulting in neuronal injury and astrocyte proliferation (astrocytosis). This post-ischemic astrocytosis is associated with up-regulated CysLT₁R and CysLT₂R [22-26]. The CysLT₁R antagonist pranlukast attenuates post-ischemic astrocytosis and glial scar formation in the chronic phases of focal cerebral ischemia in mice and rats [25,27,28]. This effect suggests that CysLT₁R mediates CysLT-induced astrocytosis and glial scar formation in response to *in vivo* ischemic injury. In primary astrocyte cultures, CysLTs are released after oxygen-glucose deprivation-induced ischemic injury, and the resultant activation of CysLT₁R mediates astrocyte proliferation [29,30]. These findings imply that the endogenously released CysLTs might play an autocrine role in the induction of astrocytosis and resultant glial scar formation through activating CysLT₁R.

However, whether CysLT₁R mediates astrocyte migration in the process of glial scar formation needs investigation. In the periphery, CysLT₁R mediates migration in many types of cells, such as monocytes [31], dendritic cells [32], monocyte-derived dendritic cells [33], vascular smooth muscle cells [34], intestinal epithelial cells [35]

and endothelial cells [31,34-36]. Therefore, CysLT₁R may also be an inducer of astrocyte migration, but many other factors have been reported to be potent inducers, such as TGF- β 1 [37,38]. Thus, there may be interactions between CysLT₁R and other regulators (for example, TGF- β 1). TGF- β 1 up-regulates CysLT₁R expression and increases the production of CysLTs in several cell types such as hepatic stellate cells [39] and bronchial smooth muscle cells [37]. Based on these findings, it is possible that the regulatory role of TGF- β 1 in astrocyte migration may be mediated by enhanced production of CysLTs *via* CysLT₁R activation. To clarify this possibility, in the present study, we investigated the interactions between TGF- β 1 and 5-LOX/CysLT₁R in astrocyte migration.

Methods

Primary cultures of rat astrocytes

Primary astrocytes were isolated from the cerebral cortex of newborn Sprague-Dawley rats within 24 h as described previously [30,40]. In brief, the cortices were digested with 0.25% trypsin and plated into poly-L-lysine-coated flasks. Cells were cultured in high-glucose DMEM (Gibco, Grand Island, NY, USA) supplemented with 10% fetal bovine serum (FBS), 2 mM glutamine, 100 units/mL penicillin and 100 μ g/mL streptomycin at 37°C in a humidified atmosphere of 95% air/5% CO₂. After incubation for 11 to 14 days, the confluent cultures were shaken overnight at 260 rpm at 37°C, and the adherent cells were trypsinized and re-seeded in the growth medium. More than 95% of the cells were astrocytes as confirmed by immunofluorescence staining for glial fibrillary acidic protein (GFAP).

All animal experiments were carried out in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. We made every effort to minimize the number of animals used and their suffering. The experimental protocols were approved by the Ethics Committee of Laboratory Animal Care and Welfare, School of Medicine, Zhejiang University.

Cell migration (wound healing) assay

Astrocytes were grown to confluence in 24-well plates and starved in serum-free DMEM for 24 h. The monolayer cells were manually scratched with a 20- μ l pipette tip to create an extended and definite scratch in the center of the dish with a bright and clear field. The detached cells were removed by washing with phosphate-buffered saline (PBS). DMEM containing 1% FBS with or without TGF- β 1 (PeproTech Inc, Rocky Hill, NJ, USA) was added to each dish. In some experiments, 1 ng/ml TGF- β 1 was added to each dish for 30 minutes before treatment with LTD₄ (Sigma-Aldrich Co., St Louis, MO, USA) or N-methyl LTC₄ (NMLTC₄, a metabolically stable LTC₄ mimetic; Cayman Chemical

Co., Ann Arbor, MI, USA). Cells were pretreated with the following inhibitor and antagonists: zileuton (0.01 to 5 μ M, a 5-LOX inhibitor; Gaomeng Pharmaceutical Co., Beijing, China), montelukast (0.01 to 5 μ M, a selective CysLT₁R antagonist; Merck & Co., Inc., Whitehouse Station, NJ, USA), and Bay cysLT₂ (0.01 to 5 μ M, a selective CysLT₂R antagonist; a kind gift from Dr. T. Jon Seiders of Amira Pharmaceuticals, Inc., San Diego, CA, USA) for 30 minutes, and then incubated with TGF- β 1 for 24 h. Images of migratory cells from the scratch boundary were acquired at 0 and 24 h under a light microscope with a digital camera.

To continuously monitor migration time-course in live astrocytes, astrocytes were plated in 35-mm dishes and grown to confluence, and then the cells were scratched and treated with LTD₄ or/and TGF- β 1 as described above. The movements of live astrocytes was traced under an inverse videomicroscope (Olympus IX81, Olympus Corp., Tokyo, Japan), and the wound was photographed at 0, 6, 12, 18 and 24 h.

The wounded areas were analyzed with ImageTool 2.0 software (University of Texas Health Science Center, San Antonio, TX, USA). The wound healing effect is determined as the initial scratch area (0 h) after wounding minus the scratch area after treatment for 24 h, or 6, 12, 18 and 24 h (live astrocytes), and reported as percentages of control values. Moreover, some astrocyte samples seeded on coverslips were visualized by GFAP


immunofluorescence staining 24 h after scratching as the typical images.

Cell proliferation assay

To measure astrocyte proliferation, carboxyfluorescein diacetate succinimidyl ester (CFSE) green fluorescent dye (Invitrogen Corp., Carlsbad, CA, USA) dilution assay was performed according to the manufacturer's instructions and the reported method [41-43]. Briefly, astrocytes were grown to confluence in six-well plates and starved in serum-free DMEM for 24 h, then the cells were washed twice with PBS and incubated in 5 μ M CFSE in PBS for 15 minutes at 37°C, and subsequently washed twice with PBS. Then DMEM containing 1% FBS with or without TGF- β 1 or LTD₄ was added to each plate. In some experiments, 1 ng/ml TGF- β 1 was added to each plate for 30 minutes before treatment with LTD₄. The cells were harvested at 24 h, and subjected to fluorescence activated cell sorting using the FC500MCL flow cytometer (Beckman Coulter, Inc., Brea, CA, USA). Proliferation was measured by loss of CFSE dye.

CysLT₁ receptor knockdown by small interfering RNA (siRNA)

RNA duplexes of 21 nucleotides specific for rat CysLT₁R sequences were chemically synthesized, together with a non-silencing negative control siRNA. The CysLT₁R siRNA sense sequence was: 5'-CAG CCU UCC AAG






Figure 2 Effect of LTD₄ on astrocyte migration and TGF-β1-induced migration and proliferation. (A, B), Photomicrographs showing astrocyte migration 24 h after treatment with LTD₄ (0.01 to 100 nmol/L) in the absence (A) or presence of TGF-β1 (1 ng/ml, B). Scale bars, 400 μm. (C) Data are reported as mean ± S.E.M.; n = 8; *P < 0.05 and **P < 0.01 compared with control, #P < 0.05 and ##P < 0.01 compared with TGF-β1 alone (LTD₄ 0). (D, E) MFI at 24 h was no significant change 24 h after treatment with LTD₄ (0.01 to 100 nM, D) alone or combined with TGF-β1 1 ng/ml (E) (n = 9 for each group, P > 0.05).

UAU ACA UTT-3' and anti-sense: 5'-AUG UAU ACU UGG AAG GCU GTT-3'; the non-silencing control siRNA sense: 5'-UUC UCC GAA CGU GUC ACG UTT-3' and anti-sense: 5'-ACG UGA CAC GUU CGG AGA ATT-3' (GenePharma Co., Shanghai, China). Transfection of siRNA duplexes was performed according to the manufacturer's instructions. Briefly, astrocytes were seeded on the day before transfection using an appropriate medium with 10% FBS without antibiotics. They were transiently transfected with CysLT₁R siRNA or negative control siRNA (100 nM) for 6 h using Lipofectamine™ 2000 (Invitrogen, USA). After the transfected cells were incubated for 48 h, they were treated with LTD₄ or TGF-β1 for cell migration assay.

Reverse transcription-polymerase chain reaction (RT-PCR)

At the end of the experiments, total RNA was extracted from the cultured astrocytes using Trizol reagent

(Invitrogen, USA) according to the manufacturer's instructions. The cDNA synthesis and PCR reactions were performed as reported previously [29,30]. The PCR primers were: 5-LOX forward 5'-AAA GAA CTG GAA ACA GCT CAG AAA-3' and reverse 5'-AAC TGG TGT GTA CAG GGG TCA GTT-3'; CysLT₁R, forward 5'- ATG TTC ACA AAG GCA AGT GG -3' and reverse 5'-TGC ATC CTA AGG ACA GAG TCA -3'; CysLT₂R, forward 5'- ACC CCT TCC AGA TGC TCC A -3' and reverse 5'- CGT GCT TTG AAA TTC TCT CCA -3'; β-actin, forward 5'-AAC CCT AAG GCC AACCGT GAA-3' and reverse 5'-TCA TGA GGT AGT CTG TCA GGT C-3'; TGF-β1, forward 5'- GAC CGC AAC AAC GCA ATC TA -3' and reverse 5'- AGG TGT TGA GCC CTT TCC AG -3'.

For cDNA synthesis, 2 μg total RNA was mixed with 1 mM deoxynucleotide triphosphate, 0.2 μg random primer, 20 U RNasin and 200 U M-MuLV reverse




Figure 3 Effect of NMLTC₄ on astrocyte migration. (A) Photomicrographs showing astrocyte migration 24 h after treatment with NMLTC₄ (0.01 to 100 nmol/L). (B) Data are reported as mean \pm S.E.M.; $n = 8$. Scale bar, 400 μ m.

transcriptase in 20 μ l reverse reaction buffer. The mixture was incubated at 42°C for 60 minutes, and then heated at 72°C for 10 minutes to inactivate the reverse transcriptase.

PCR was performed on an Eppendorf Master Cycler (Eppendorf Scientific, Inc., Westbury, NY, USA) as follows: 1 μ l cDNA mixture was reacted in 20 μ l reaction buffer containing 1.5 mM MgCl₂, 0.2 mM deoxynucleotide triphosphate, 20 pM primer and 1 U Taq DNA polymerase. The reaction mixtures were initially heated at 94°C for 2 minutes, then at 94°C for 60 sec, 56°C for 60 sec, and 72°C for 60 sec for 35 cycles and finally stopped at 72°C for 10 minutes. With the exception of TGF- β 1, the reaction mixtures were initially heated at 94°C for 2 minutes, then at 94°C for 30 sec, 54°C for 30 sec, and 72°C for 60 sec for 28 cycles and finally stopped at 72°C for 10 minutes. PCR products of 20 μ l were separated by 2% agarose gel electrophoresis and visualized by ethidium bromide staining. The density of each band was measured by a UVP gel analysis system (Bio-Rad Laboratories, Hercules, CA, USA). The results are expressed as the ratios to β -actin.

Western blotting analysis

Astrocytes were washed twice with ice-cold PBS and then lysed for 30 minutes on ice in Cell and Tissue Protein

Extraction Solution (Kangcheng Biotechnology Inc., Shanghai, China). The homogenate was centrifuged at 12,000 g for 30 minutes at 4°C, and the supernatant was used. The protein samples (100 μ g) were separated by 10% SDS-polyacrylamide gels and then transferred to nitrocellulose membranes (Invitrogen). The membranes were blocked by 10% fat-free milk, and sequentially incubated with the following antibodies: rabbit polyclonal antibody against CysLT₁R (1:200) [44], CysLT₂R (1:200) [26,45] or 5-LOX (1:300, (Chemicon International Inc. Temecula, CA, USA) and mouse monoclonal antibody against glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (1:5,000, Kangcheng Biotechnology Inc., Shanghai, China) at 4°C overnight. After repeated wash, the membranes were incubated with anti-rabbit IRDye700DX[®]-conjugated antibody or anti-mouse IRDye800DX[®]-conjugated antibody (1:5,000, Rockland Immunochemicals, Inc., Gilbertsville, PA, USA). The immunoblot was analyzed by the Odyssey Fluorescence Scanner (LI-COR Bioscience, Inc., Lincoln, NE, USA). The protein bands were quantified using BIORAD Quantity One software (Bio-Rad, USA). The results are expressed as the ratios to GAPDH.

Immunofluorescence staining

Astrocytes seeded on coverslips were fixed in cold methanol for 5 minutes, and incubated in 10% normal goat serum for 2 h to block non-specific binding of IgG. Then the cells were reacted with a mouse monoclonal antibody against GFAP (1:500, Millipore Corp., Bedford, MA, USA) and a rabbit polyclonal antibody against CysLT₁R (1:200, Chemicon, USA) at 4°C overnight. After washing in PBS, astrocytes were incubated with FITC-conjugated goat anti-mouse or Cy3-conjugated goat anti-rabbit antibody (1:200, Millipore, USA) for 2 h at room temperature. Finally, the stained cells were observed under a fluorescence microscope (Olympus BX51, Olympus Corp., Tokyo, Japan). Control coverslips were treated with normal goat serum instead of the primary antibody, and did not show positive immunostaining (data not shown).

5-LOX immunocytochemistry

Astrocytes cultured on coverslips were fixed in cold methanol (-20°C) for 5 minutes and incubated for 30 minutes in PBS containing 3% H₂O₂ to eliminate endogenous peroxidase activity. Then, cells were incubated for 2 h in PBS containing 10% normal goat serum and incubated at 4°C overnight with rabbit polyclonal antibody against 5-LOX (1:200, Chemicon, USA) as the primary antibody. After three washes with PBS, cells were incubated for 2 h with biotin-conjugated goat anti-rabbit IgG antiserum (1:200) as a second antibody, followed by incubation with avidin-biotin-HRP complex. Finally, the cells were visualized with 0.01% 3, 3'-diaminobenzidine




Figure 4 Time-dependent migration of live astrocytes after exposure to TGF-β1 and LTD₄. Live astrocytes were continuously monitored under a videomicroscope after exposure to TGF-β1 or/and LTD₄. (A) Representative images showing astrocyte migration traced by videomicroscopy at 6, 12, 18 and 24 h after scratching. Scale bar, 200 μm. (B) TGF-β1 and LTD₄ concentration- and time-dependently accelerated migration. When TGF-β1 (1 ng/ml) combined with LTD₄ (0.1 nM), the effect at 24 h was more potent than that of TGF-β1 or LTD₄ alone. Data are reported as mean ± S.E.M.; n = 3; *P < 0.05 and **P < 0.01 compared with corresponding time-points of the control; ##P < 0.01 compared with TGF-β1 + LTD₄ at 24 h.

and 0.005% H₂O₂ in 50 mM Tris-HCl, pH 7.6. Control coverslips were treated with normal goat serum instead of the primary antibody and they did not show positive immunostaining (data not shown). Then, the cells were examined under the Olympus microscope.

Measurement of extracellular cysteinyl leukotrienes and TGF-β1

According to the reported method [29,30], astrocytes were seeded into six-well culture plates at 5×10^5 cells/well in 2 ml standard culture medium for 24 h. After culture in DMEM without serum for another 24 h, astrocytes were cultured in DMEM with 1% FBS and stimulated with TGF-β1 (10 ng/mL), various concentrations of LTD₄ or NMLTC₄, or vehicle for the designated times. Then, cell-free supernatants were stored at -80°C. The CysLTs (LTC₄, LTD₄ and LTE₄) in astrocyte supernatants were assayed using a commercial CysLT ELISA kit (Cayman Chemical Co., Ann Arbor, MI, USA) according to the manufacturer's instructions and

calculated as pg/mg protein. The TGF-β1 in the supernatants was assayed using a commercial TGF-β1 ELISA kit (Wuhan Boster Biological Technology Co., Ltd., Wuhan, China) according to the manufacturer's instructions, and calculated as pg/ml.

Statistical analysis

Data are reported as mean ± S.E.M. Student's *t*-test and one-way analysis of variance were used to determine the statistical significance of differences between groups. A value of *P* < 0.05 was considered statistically significant.

Results

TGF-β1- and LTD₄-induced astrocyte migration

First, we confirmed the effect of TGF-β1 on astrocyte migration. TGF-β1 (1 and 10 ng/ml for 24 h) significantly accelerated the migration of astrocytes from the wound edge into the central area in a concentration-dependent manner (Figure 1A). To distinguish the effects on migration and proliferation, we determined whether TGF-β1




Figure 5 Effects of a 5-LOX inhibitor and CysLT receptor antagonists on TGF-β1-induced migration in astrocytes. (A) Photomicrographs showing TGF-β1 (10 ng/ml)-induced astrocyte migration 24 h after treatment with the 5-LOX inhibitor zileuton, the CysLT₁R antagonist montelukast and the CysLT₂R antagonist Bay cysLT2 (1 μM). Scale bar, 400 μm. (B-D) TGF-β1-induced migration inhibited by 0.01 to 5 μM zileuton (B) and montelukast (C), but not Bay cysLT2 (D). Data are reported as mean ± S.E.M.; n = 8; *P < 0.05 and **P < 0.01 compared with control; #P < 0.05 compared with TGF-β1 alone.

affects astrocyte proliferation. The results of CFSE fluorescence intensity showed that astrocyte proliferation did not differ from control level 24 h after exposure to TGF-β1 (0.1, 1 and 10 ng/ml) (Figure 1C) although the assay confirmed astrocyte proliferation at 24 h compared with 0 h (baseline) (Figure 1B).

Next, we determined whether the non-selective agonist LTD₄ and the CysLT₂R agonist NMLTC₄ [46] induce astrocyte migration, and LTD₄ potentiates the TGF-β1 effect. The results showed that LTD₄ significantly stimulated the migration of astrocytes at 0.1 to 10 nM but not at 0.01 and 100 nM; the maximum migration (141.7 ± 5.0%) was induced by 1 nM LTD₄ (Figure 2A, C). LTD₄ (0.01 to 1 nM) also potentiated the effect of the lower concentration of TGF-β1 (1 ng/ml); the migration rates after treatment with 1 ng/ml TGF-β1 were increased from 110.3 ± 5.4% to 175.3 ± 4.8% with 0.01 nM, from 123.5 ± 4.0% to 203.5 ± 5.3% with 0.1 nM, and from 141.7 ± 5.0% to 193.8 ± 2.9% with 1 nM LTD₄ (Figure 2B, C). LTD₄ (0.01 to 100 nM) alone (Figure 2D) or combined with TGF-β1 1 ng/ml (Figure 2E) did not affect astrocyte proliferation at 24 h. However, NMLTC₄ (0.01 to 100 nM for 24 h) did not have any significant effect on astrocyte migration (Figure 3). In addition, to confirm the migration and determine its temporal property, we continuously monitored migration of live astrocytes during 24 h after exposure to LTD₄ or/and TGF-β1. We found that TGF-β1 (1 and 10 ng/ml) and LTD₄ (1 nM) gradually accelerated migration during 24 h in a concentration-dependent manner. When TGF-β1 (1 ng/ml) combined with LTD₄ (0.1 nM), the effect at

24 h was more potent than that of TGF-β1 or LTD₄ alone (Figure 4).

To confirm the roles of endogenous CysLTs and CysLT₁R in TGF-β1-induced migration, we examined the effects of the 5-LOX inhibitor zileuton, the CysLT₁R antagonist montelukast, and the CysLT₂R antagonist Bay cysLT2 as well as CysLT₁R siRNA. We found that the effect of 10 ng/ml TGF-β1 was attenuated by zileuton (1 and 5 μM, Figure 5A, B) and montelukast (1 and 5 μM, Figure 5A, C), but not by Bay cysLT2 (0.01 to 5 μM, Figure 5A, D). These results indicated that endogenously released CysLTs might activate CysLT₁R, but not CysLT₂R, to induce astrocyte migration and potentiate TGF-β1-induced migration. The involvement of CysLT₁R was further confirmed by RNA silencing by transient transfection of CysLT₁R siRNA into astrocytes. The siRNA (100 nM) significantly reduced the expression of CysLT₁R mRNA (Figure 6A) and protein (Figure 6B, C), but the non-silencing negative control siRNA had no effect. CysLT₁R siRNA significantly attenuated the effects of LTD₄ (1 and 10 nM) and TGF-β1 (1 and 10 ng/ml) on astrocyte migration (Figure 6D, E). These results suggest that CysLT₁R may be associated with LTD₄- and TGF-β1-induced astrocyte migration.

TGF-β1-Induced Activation of 5-LOX in astrocytes

To investigate the role of endogenous CysLTs, the 5-LOX metabolites, in TGF-β1-induced astrocyte migration, we determined 5-LOX expression in astrocytes. We found that TGF-β1 10 ng/ml significantly increased 5-LOX mRNA (Figure 7A) and protein expression




Figure 6 Effect of CysLT₁R siRNA on LTD₄- and TGF-β1-induced migration in astrocytes. (A, B) RT-PCR and Western blotting results showing inhibition of CysLT₁R, but not CysLT₂R, mRNA (A) and protein expression (B) by CysLT₁R siRNA but not by negative control (NC) siRNA. (C) Double immunofluorescence staining showing inhibition of CysLT₁R protein expression by CysLT₁R siRNA in GFAP-positive astrocytes. (D) Photomicrographs showing that the astrocyte migration induced by LTD₄ (1 and 10 nM) and TGF-β1 (1 and 10 ng/ml) was inhibited by CysLT₁R siRNA (siRNA) but not by NC siRNA. (E) CysLT₁R siRNA inhibited migration induced by LTD₄ and TGF-β1. Data are reported as mean ± S.E.M.; n = 4 (A and B) or 8 (E); *P < 0.05 and **P < 0.01 compared with control; #P < 0.05 and ##P < 0.01 compared with LTD₄ or TGF-β1 alone. Scale bar, 200 μm (C) or 400 μm (D).

(Figure 7B) 24 h after exposure. Immunocytochemical results showed that 5-LOX was translocated from the cytosol to the nuclear envelope 6 and 12 h after exposure to 10 ng/ml TGF-β1, and then recovered at 24 h (Figure 7C). We further determined the changes in enzymatic activity of 5-LOX by measuring its metabolites, CysLTs, in the culture medium. The levels of CysLTs increased from 1.5 h, peaked at 12 h, and were sustained over 24 h after exposure to 10 ng/ml TGF-β1 (Figure 7D). These findings revealed the involvement of 5-LOX and its metabolite CysLTs in the responses to TGF-β1.


TGF-β1-regulated expression of CysLT receptor in astrocytes

Finally, we determined whether TGF-β1 regulates the expression of CysLT₁R and CysLT₂R mRNA and protein in astrocytes, and whether LTD₄ regulates TGF-β1 expression and release. RT-PCR and Western blot showed weak expression of CysLT₁R and CysLT₂R in control astrocytes. Exposure to 10 ng/ml TGF-β1 for 24 h


induced about three-fold increase in the mRNA (Figure 8A) and protein expression (Figure 8B) of CysLT₁R, but did not significantly change the expression of CysLT₂R. Immunofluorescence staining confirmed the enhancement of CysLT₁R by TGF-β1 (Figure 8C). On the other hand, treatment with various concentrations of LTD₄ or NMLTC₄ for 24 h did not affect the TGF-β1 mRNA expression in astrocytes (Figure 9A) and its content in the culture medium (Figure 9B). Thus, TGF-β1 might up-regulate CysLT₁R but is not regulated by LTD₄.

Discussion

In the present study, we revealed that TGF-β1-induced astrocyte migration is, at least in part, mediated by enhanced endogenous CysLTs through activation of CysLT₁R. The evidence is that TGF-β1-induced astrocyte migration was potentiated by LTD₄ but attenuated by a 5-LOX inhibitor and a CysLT₁R antagonist, and TGF-β1 activated 5-LOX and increased CysLT₁R




expression. Our observations have confirmed the TGF-β1-induced migration of rat astrocytes as reported [6], and indicated another mechanism underlying TGF-β1-induced astrocyte migration in addition to the pathways through activation of the Smad family [47,48] or the ROS-dependent ERK/JNK-NF-κB pathway [6]. In addition, we found that both TGF-β1 and LTD₄ did not alter astrocyte proliferation during 24 h. It has been reported that TGF-β1 inhibits astrocyte proliferation [47,49,50] and LTD₄ induces the proliferation via activating CysLT₁R [30]. This difference between these reported results and ours may result from different assessment timing [30] and methods [47,49,50]. However, in our experimental conditions, TGF-β1 and LTD₄ regulate astrocyte migration rather than proliferation.




TGF-β1-induced astrocyte migration might be mediated by the CysLT signal pathway in at least two ways, that is, TGF-β1 potentiates the activity of both 5-LOX and CysLT₁R. On one hand, TGF-β1 increased 5-LOX expression and induced its translocation to the nuclear envelope (Figure 7C), a key step for 5-LOX activation [51-53] and, thereby, increased the production of endogenous CysLTs (Figure 7D). Consistent with this, it has been reported that TGF-β1 induces 5-LOX expression in myeloid cell lines [54-58]. The notion is also supported by the finding that the TGF-β1 effect was inhibited by the 5-LOX inhibitor zileuton (Figure 5A). On the other hand, TGF-β1 potentiates the expression of CysLT₁R, enhancing the activity of endogenously-produced or exogenous CysLTs as previously reported [37,39]. Therefore, one of the mechanisms underlying TGF-β1-induced astrocyte migration may be activation of endogenous 5-LOX /CysLT₁R signals.

Here, we demonstrated that the receptor subtype that mediated the TGF-β1 effect was CysLT₁R. The evidence was from the different effects of agonists and antagonists, and the effect of RNA interference. The non-selective agonist LTD₄ induced a moderate migration of astrocytes at lower concentrations (0.1 to 10 nM), but not at the higher concentrations 100 nM (Figure 2A, C) and 1,000 nM (data not shown). This concentration-response relationship indicated that CysLT₁R might



mediate the effect of LTD₄, because CysLT₁R is activated at 1 to 10 nM while CysLT₂R is activated at 100 to 1,000 nM in astrocytes [30]. This is also supported by the finding that the selective CysLT₂R agonist NMLTC₄ [46] had no effect on astrocyte migration (Figure 3). With regard



to receptor antagonism, the effect of TGF-β1 was attenuated by the CysLT₁R antagonist montelukast but not by the CysLT₂R antagonist Bay cysLT2. Bay cysLT2 is at least 100- to 500-fold more selective for CysLT₂R versus CysLT₁R; its pA₂ value indicates that at least 5 μM would act on the CysLT₁R [59,60]. Thus, lacking the effect of 5 μM Bay ctsLT2 in our study may be due to cell specificity and response difference. On the other hand, interference with CysLT₁R siRNA inhibited both TGF-β1- and LTD₄-induced astrocyte migration by down-regulating the expression of this receptor (Figure 6). These findings are consistent with reports that CysLT₁R mediates the migration of other types of cells [31-36]. Therefore, CysLT₁R is an important regulator of astrocyte migration in addition to its regulation of astrocyte proliferation [29,30].

The interaction between TGF-β1 and CysLTs was also investigated by determining the action of LTD₄ or NMLTC₄ on TGF-β1 expression and release. Unlike the action of TGF-β1 on the production of CysLTs and LTD₄ effects, LTD₄ or NMLTC₄ affected neither TGF-β1 expression nor its release in astrocytes (Figure 9). This may depend on specific cell types because LTD₄ induces TGF-β1 mRNA expression in human bronchial epithelial cells [61,62] and in fibroblasts from asthmatics [63], and LTC₄ induces TGF-β1 production in airway epithelium [62] in a CysLT₁R-dependent manner. Anyway, the effect of LTD₄ on TGF-β1 in astrocytes remains to be further investigated, especially in animal models of chronic brain injury. Since both levels of TGF-β1 and CysLTs are increased after brain injury [24,64,65] and involved in glial scar formation [25,65,66]; which of them is determinant in glial scar formation should be clarified for their therapeutic implications. Herein, our results suggest that activation of the endogenous 5-LOX/CysLT₁R signals might be an intermediate event in TGF-β1-regulated astrocyte migration, but not the initial event. Since TGF-β1 signaling is mainly modulated by Smad-dependent [67-75] and Smad-independent pathways [6,76-81], whether the regulation mode is mediated by the Smad or other pathways requires investigation.

Astrocyte migration is a critical step in the formation of a densely-packed glial scar [1,2], and TGF-β1 is closely associated with glial scar formation [64,66,82-84]. Thus, CysLT receptor antagonists or 5-LOX inhibitors may be beneficial in the prevention and attenuation of glial scar formation after brain injury. Actually, we have reported that the CysLT₁R antagonist pranlukast attenuates glial scar formation in the chronic phase of focal cerebral ischemia in mice [28] and rats [25], and the 5-LOX inhibitor caffeic acid has this effect in rats with focal cerebral ischemia [85] and in mice with brain cryoinjury [86]. Moreover, montelukast inhibits the astrocyte proliferation induced by mild ischemia-like

injury and low concentrations of LTD₄ [30]. The present study highlights the previous findings and clarifies the mode of action of endogenous CysLTs/CysLT₁R in the critical step of glial scar formation.

In conclusion, in the present study we found that TGF- β 1-induced astrocyte migration is, at least in part, mediated by enhanced endogenous CysLTs through activating up-regulated CysLT₁R (Figure 10). These findings indicate that the interaction between the cytokine TGF- β 1 and pro-inflammatory mediators (CysLTs) are involved in the regulation of astrocyte function relevant to glial scar formation. However, the detailed mechanisms underlying this interaction need investigation.

Abbreviations

5-LOX: 5-lipoxygenase; CNS: central nervous system; CysLT₁R: cysteinyl leukotriene receptor 1; CysLT₂R: cysteinyl leukotriene receptor 2; CysLTs: Cysteinyl leukotrienes; FBS: fetal bovine serum; GAPDH: glyceraldehyde 3-phosphate dehydrogenase; GFAP: glial fibrillary acidic protein; LTD₄: leukotriene D₄; MFI: mean fluorescence intensity; NMLTC₄: N-methyl leukotriene C₄; PBS: phosphate-buffered saline; siRNA: small interfering RNA; TGF- β 1: transforming growth factor- β 1.

Competing interests

The authors have no competing interests.

Acknowledgments

We thank Dr. T. Jon Seiders, Amira Pharmaceuticals Inc., USA, for supplying BAY cysLT₂; Dr. John Obenhain, Merck Research Laboratories, USA, for montelukast; and Dr. IC Bruce for critically reading and revising this manuscript. This study was supported by the National Natural Science Foundation of China (81072618, 81173041, 30772561 and 30873053), the Zhejiang Provincial Natural Science Foundation (Y2090069), the Zhejiang Provincial "QianjiangRenCai Research Plan" (2010R10055), and the Fundamental Research Funds for the Central Universities (2009QNA7008).

Authors' contributions

XQH designed the study, performed the main parts of the experiments, analyzed the data, and prepared the manuscript. XYZ performed the immunofluorescence staining experiments and prepared the figures. XRW performed the cell migration experiments and analyzed the data. YYY performed the RT-PCR experiments and analyzed the data. SHF, YBL and WPZ contributed to the design of the study, to interpretation of the results and to the writing of the manuscript. EQW made essential contributions to the design of the study and interpretation of the results, and completed the manuscript. All authors read and approved the final manuscript.

Received: 14 January 2012 Accepted: 17 May 2012

Published: 26 June 2012

References

1. Fawcett JW, Asher RA: **The glial scar and central nervous system repair.** *Brain Res Bull* 1999, **49**:377–391.
2. Saadoun S, Papadopoulos MC, Watanabe H, Yan D, Manley GT, Verkman AS: **Involvement of aquaporin-4 in astroglial cell migration and glial scar formation.** *J Cell Sci* 2005, **118**:5691–5698.
3. Faber-Elman A, Lavie V, Schwartz I, Shaltiel S, Schwartz M: **Vitronectin overrides a negative effect of TNF- α on astrocyte migration.** *FASEB J* 1995, **9**:1605–1613.
4. Striedinger K, Scemes E: **Interleukin-1 β affects calcium signaling and *in vitro* cell migration of astrocyte progenitors.** *J Neuroimmunol* 2008, **196**:116–123.
5. Miao H, Crabb AW, Hernandez MR, Lukas TJ: **Modulation of factors affecting optic nerve head astrocyte migration.** *Invest Ophthalmol Vis Sci* 2010, **51**:4096–4103.
6. Hsieh HL, Wang HH, Wu WB, Chu PJ, Yang CM: **Transforming growth factor- β 1 induces matrix metalloproteinase-9 and cell migration in**

- astrocytes: roles of ROS-dependent ERK- and JNK-NF- κ B pathways. *J Neuroinflammation* 2010, **7**:88.
7. Flanders KC, Ren RF, Lippa CF: **Transforming growth factor- β in neurodegenerative disease.** *Prog Neurobiol* 1998, **54**:71–85.
8. Unsicker K, Strelau J: **Functions of transforming growth factor- β isoforms in the nervous system. Cues based on localization and experimental *in vitro* and *in vivo* evidence.** *Eur J Biochem* 2000, **267**:6972–6975.
9. Bottner M, Kriegstein K, Unsicker K: **The transforming growth factor- β s: structure, signaling, and roles in nervous system development and functions.** *J Neurochem* 2000, **75**:2227–2240.
10. Massague J: **How cells read TGF- β signals.** *Nat Rev Mol Cell Biol* 2000, **1**:169–178.
11. Vivien D, Ali C: **Transforming growth factor- β signalling in brain disorders.** *Cytokine Growth Factor Rev* 2006, **17**:121–128.
12. Leivonen SK, Kahari VM: **Transforming growth factor- β signaling in cancer invasion and metastasis.** *Int J Cancer* 2007, **121**:2119–2124.
13. Pratt BM, McPherson JM: **TGF- β in the central nervous system: potential roles in ischemic injury and neurodegenerative diseases.** *Cytokine Growth Factor Rev* 1997, **8**:267–292.
14. Lehmann E, Kiefer R, Christensen T, Toyka KV, Zimmer J, Diemer NH, Hartung HP, Finsen B: **Microglia and macrophages are major sources of locally produced transforming growth factor- β 1 after transient middle cerebral artery occlusion in rats.** *Glia* 1998, **24**:437–448.
15. Ruocco A, Nicole O, Docagne F, Ali C, Chazalviel L, Komesli S, Yablonsky F, Roussel S, MacKenzie ET, Vivien D, Buisson A: **A transforming growth factor- β antagonist unmasks the neuroprotective role of this endogenous cytokine in excitotoxic and ischemic brain injury.** *J Cereb Blood Flow Metab* 1999, **19**:1345–1353.
16. Samuelsson B, Dahlen SE, Lindgren JA, Rouzer CA, Serhan CN: **Leukotrienes and lipoxins: structures, biosynthesis, and biological effects.** *Science* 1987, **237**:1171–1176.
17. Kanaoka Y, Boyce JA: **Cysteinyl leukotrienes and their receptors: cellular distribution and function in immune and inflammatory responses.** *J Immunol* 2004, **173**:1503–1510.
18. Wenzel SE: **The role of leukotrienes in asthma.** *Prostaglandins Leukot Essent Fatty Acids* 2003, **69**:145–155.
19. Vannella KM, McMillan TR, Charbeneau RP, Wilke CA, Thomas PE, Toews GB, Peters-Golden M, Moore BB: **Cysteinyl leukotrienes are autocrine and paracrine regulators of fibrocyte function.** *J Immunol* 2007, **179**:7883–7890.
20. Brink C, Dahlen SE, Drazen J, Evans JF, Hay DW, Nicosia S, Serhan CN, Shimizu T, Yokomizo T: **International union of pharmacology XXXVII. Nomenclature for leukotriene and lipoxin receptors.** *Pharmacol Rev* 2003, **55**:195–227.
21. Rovati GE, Capra V: **Cysteinyl-leukotriene receptors and cellular signals.** *Scientific World Journal* 2007, **7**:1375–1392.
22. Fang SH, Zhou Y, Chu LS, Zhang WP, Wang ML, Yu GL, Peng F, Wei EQ: **Spatio-temporal expression of cysteinyl leukotriene receptor-2 mRNA in rat brain after focal cerebral ischemia.** *Neurosci Lett* 2007, **412**:78–83.
23. Zhang YJ, Zhang L, Ye YL, Fang SH, Zhou Y, Zhang WP, Lu YB, Wei EQ: **Cysteinyl leukotriene receptors CysLT1 and CysLT2 are upregulated in acute neuronal injury after focal cerebral ischemia in mice.** *Acta Pharmacol Sin* 2006, **27**:1553–1560.
24. Zhou Y, Wei EQ, Fang SH, Chu LS, Wang ML, Zhang WP, Yu GL, Ye YL, Lin SC, Chen Z: **Spatio-temporal properties of 5-lipoxygenase expression and activation in the brain after focal cerebral ischemia in rats.** *Life Sci* 2006, **79**:1645–1656.
25. Fang SH, Wei EQ, Zhou Y, Wang ML, Zhang WP, Yu GL, Chu LS, Chen Z: **Increased expression of cysteinyl leukotriene receptor-1 in the brain mediates neuronal damage and astrogliosis after focal cerebral ischemia in rats.** *Neuroscience* 2006, **140**:969–979.
26. Zhao CZ, Zhao B, Zhang XY, Huang XQ, Shi WZ, Liu HL, Fang SH, Lu YB, Zhang WP, Tang FD, Wei EQ: **Cysteinyl leukotriene receptor 2 is spatiotemporally involved in neuron injury, astrogliosis and microgliosis after focal cerebral ischemia in rats.** *Neuroscience* 2011, **189**:1–11.
27. Yu GL, Wei EQ, Zhang SH, Xu HM, Chu LS, Zhang WP, Zhang Q, Chen Z, Mei RH, Zhao MH: **Montelukast, a cysteinyl leukotriene receptor-1 antagonist, dose- and time-dependently protects against focal cerebral ischemia in mice.** *Pharmacology* 2005, **73**:31–40.
28. Yu GL, Wei EQ, Wang ML, Zhang WP, Zhang SH, Weng JQ, Chu LS, Fang SH, Zhou Y, Chen Z, Zhang Q, Zhang LH: **Pranlukast, a cysteinyl leukotriene**

- receptor-1 antagonist, protects against chronic ischemic brain injury and inhibits the glial scar formation in mice. *Brain Res* 2005, **1053**:116–125.
29. Ciccarelli R, D'Alimonte I, Santavenero C, D'Auro M, Ballerini P, Nargi E, Buccella S, Nicosia S, Folco G, Caciagli F, Di Lorio P: **Cysteinyl-leukotrienes are released from astrocytes and increase astrocyte proliferation and glial fibrillary acidic protein via cys-LT1 receptors and mitogen-activated protein kinase pathway.** *Eur J Neurosci* 2004, **20**:1514–1524.
30. Huang XJ, Zhang WP, Li CT, Shi WZ, Fang SH, Lu YB, Chen Z, Wei EQ: **Activation of CysLT receptors induces astrocyte proliferation and death after oxygen-glucose deprivation.** *Glia* 2008, **56**:27–37.
31. Woszczek G, Chen LY, Nagineni S, Kern S, Barb J, Munson PJ, Logun C, Danner RL, Shelhamer JH: **Leukotriene D(4) induces gene expression in human monocytes through cysteinyl leukotriene type I receptor.** *J Allergy Clin Immunol* 2008, **121**:215–221. e211.
32. Thivierge M, Stankova J, Rola-Pleszczynski M: **Toll-like receptor agonists differentially regulate cysteinyl-leukotriene receptor 1 expression and function in human dendritic cells.** *J Allergy Clin Immunol* 2006, **117**:1155–1162.
33. Thivierge M, Stankova J, Rola-Pleszczynski M: **Cysteinyl-leukotriene receptor type 1 expression and function is down-regulated during monocyte-derived dendritic cell maturation with zymosan: involvement of IL-10 and prostaglandins.** *J Immunol* 2009, **183**:6778–6787.
34. Kaetsu Y, Yamamoto Y, Sugihara S, Matsuura T, Igawa G, Matsubara K, Igawa O, Shigemasa C, Hisatome I: **Role of cysteinyl leukotrienes in the proliferation and the migration of murine vascular smooth muscle cells in vivo and in vitro.** *Cardiovasc Res* 2007, **76**:160–166.
35. Paruchuri S, Broom O, Dib K, Sjolander A: **The pro-inflammatory mediator leukotriene D4 induces phosphatidylinositol 3-kinase and Rac-dependent migration of intestinal epithelial cells.** *J Biol Chem* 2005, **280**:13538–13544.
36. Yuan YM, Fang SH, Qian XD, Liu LY, Xu LH, Shi WZ, Zhang LH, Lu YB, Zhang WP, Wei EQ: **Leukotriene D4 stimulates the migration but not proliferation of endothelial cells mediated by the cysteinyl leukotriene cyslt(1) receptor via the extracellular signal-regulated kinase pathway.** *J Pharmacol Sci* 2009, **109**:285–292.
37. Espinosa K, Bosse Y, Stankova J, Rola-Pleszczynski M: **CysLT1 receptor upregulation by TGF-beta and IL-13 is associated with bronchial smooth muscle cell proliferation in response to LTD4.** *J Allergy Clin Immunol* 2003, **111**:1032–1040.
38. Asakura T, Ishii Y, Chibana K, Fukuda T: **Leukotriene D4 stimulates collagen production from myofibroblasts transformed by TGF-beta.** *J Allergy Clin Immunol* 2004, **114**:310–315.
39. Paiva LA, Maya-Monteiro CM, Bandeira-Melo C, Silva PM, El-Cheikh MC, Teodoro AJ, Borojevic R, Perez SA, Bozza PT: **Interplay of cysteinyl leukotrienes and TGF-beta in the activation of hepatic stellate cells from Schistosoma mansoni granulomas.** *Biochim Biophys Acta* 2010, **1801**:1341–1348.
40. Qi LL, Fang SH, Shi WZ, Huang XQ, Zhang XY, Lu YB, Zhang WP, Wei EQ: **CysLT2 receptor-mediated AQP4 up-regulation is involved in ischemic-like injury through activation of ERK and p38 MAPK in rat astrocytes.** *Life Sci* 2011, **88**:50–56.
41. Deleyrolle LP, Harding A, Cato K, Siebzehrubel FA, Rahman M, Azari H, Olson S, Gabrielli B, Osborne G, Vescovi A, Reynolds BA: **Evidence for label-retaining tumour-initiating cells in human glioblastoma.** *Brain* 2011, **134**:1331–1343.
42. Quah BJ, Parish CR: **New and improved methods for measuring lymphocyte proliferation in vitro and in vivo using CFSE-like fluorescent dyes.** *J Immunol Methods* 2012, **379**:1–14.
43. Bogie JF, Stinissen P, Hellings N, Hendriks JJ: **Myelin-phagocytosing macrophages modulate autoreactive T cell proliferation.** *J Neuroinflammation* 2011, **8**:85.
44. Luo JY, Zhang Z, Yu SY, Zhao B, Zhao CZ, Wang XX, Fang SH, Zhang WP, Zhang LH, Wei EQ, Lu YB: **Rotenone-induced changes of cysteinyl leukotriene receptor 1 expression in BV2 microglial cells.** *Zhejiang Da Xue Xue Bao Yi Xue Ban* 2011, **40**:131–138.
45. Zhang LP, Zhao CZ, Shi WZ, Qi LL, Lu YB, Zhang YM, Zhang LH, Fang SH, Bao JF, Shen JG, Wei EQ: **Preparation and identification of polyclonal antibody against cysteinyl leukotriene receptor 2.** *Zhejiang Da Xue Xue Bao Yi Xue Ban* 2009, **38**:591–597.
46. Yan D, Stocco R, Sawyer N, Nesheim ME, Abramovitz M, Funk CD: **Differential signaling of cysteinyl leukotrienes and a novel cysteinyl leukotriene receptor 2 (CysLT) agonist, N-methyl-leukotriene C, in calcium reporter and beta arrestin assays.** *Mol Pharmacol* 2011, **79**:270–278.
47. Stipursky J, Gomes FC: **TGF-beta1/SMAD signaling induces astrocyte fate commitment in vitro: implications for radial glia development.** *Glia* 2007, **55**:1023–1033.
48. Schachtrup C, Ryu JK, Helmrick MJ, Vagena E, Galanakis DK, Degen JL, Margolis RU, Akassoglou K: **Fibrinogen triggers astrocyte scar formation by promoting the availability of active TGF-beta after vascular damage.** *J Neurosci* 2010, **30**:5843–5854.
49. Lindholm D, Castren E, Kiefer R, Zafra F, Thoenen H: **Transforming growth factor-beta 1 in the rat brain: increase after injury and inhibition of astrocyte proliferation.** *J Cell Biol* 1992, **117**:395–400.
50. Klaver CL, Caplan MR: **Bioactive surface for neural electrodes: decreasing astrocyte proliferation via transforming growth factor-beta1.** *J Biomed Mater Res A* 2007, **81**:1011–1016.
51. Ge QF, Wei EQ, Zhang WP, Hu X, Huang XJ, Zhang L, Song Y, Ma ZQ, Chen Z, Luo JH: **Activation of 5-lipoxygenase after oxygen-glucose deprivation is partly mediated via NMDA receptor in rat cortical neurons.** *J Neurochem* 2006, **97**:992–1004.
52. Li CT, Zhang WP, Lu YB, Fang SH, Yuan YM, Qi LL, Zhang LH, Huang XJ, Zhang L, Chen Z, Wei EQ: **Oxygen-glucose deprivation activates 5-lipoxygenase mediated by oxidative stress through the p38 mitogen-activated protein kinase pathway in PC12 cells.** *J Neurosci Res* 2009, **87**:991–1001.
53. Song Y, Wei EQ, Zhang WP, Ge QF, Liu JR, Wang ML, Huang XJ, Hu X, Chen Z: **Minocycline protects PC12 cells against NMDA-induced injury via inhibiting 5-lipoxygenase activation.** *Brain Res* 2006, **1085**:57–67.
54. Steinhilber D, Radmark O, Samuelsson B: **Transforming growth factor beta upregulates 5-lipoxygenase activity during myeloid cell maturation.** *Proc Natl Acad Sci USA* 1993, **90**:5984–5988.
55. Brungs M, Radmark O, Samuelsson B, Steinhilber D: **On the induction of 5-lipoxygenase expression and activity in HL-60 cells: effects of vitamin D3, retinoic acid, DMSO and TGF beta.** *Biochem Biophys Res Commun* 1994, **205**:1572–1580.
56. Brungs M, Radmark O, Samuelsson B, Steinhilber D: **Sequential induction of 5-lipoxygenase gene expression and activity in Mono Mac 6 cells by transforming growth factor beta and 1,25-dihydroxyvitamin D3.** *Proc Natl Acad Sci USA* 1995, **92**:107–111.
57. Harle D, Radmark O, Samuelsson B, Steinhilber D: **Calcitriol and transforming growth factor-beta upregulate 5-lipoxygenase mRNA expression by increasing gene transcription and mRNA maturation.** *Eur J Biochem* 1998, **254**:275–281.
58. Seuter S, Sorg BL, Steinhilber D: **The coding sequence mediates induction of 5-lipoxygenase expression by Smads3/4.** *Biochem Biophys Res Commun* 2006, **348**:1403–1410.
59. Ni NC, Yan D, Ballantyne LL, Barajas-Espinosa A, St Amand T, Pratt DA, Funk CD: **A selective cysteinyl leukotriene receptor 2 antagonist blocks myocardial ischemia/reperfusion injury and vascular permeability in mice.** *J Pharmacol Exp Ther* 2011, **339**:768–778.
60. Carnini C, Accomazzo MR, Borroni E, Vitellaro-Zuccarello L, Durand T, Folco G, Rovati GE, Capra V, Sala A: **Synthesis of cysteinyl leukotrienes in human endothelial cells: subcellular localization and autocrine signaling through the CysLT2 receptor.** *FASEB J* 2011, **25**:3519–3528.
61. Bosse Y, Thompson C, McMahon S, Dubois CM, Stankova J, Rola-Pleszczynski M: **Leukotriene D4-induced, epithelial cell-derived transforming growth factor beta1 in human bronchial smooth muscle cell proliferation.** *Clin Exp Allergy* 2008, **38**:113–121.
62. Perng DW, Wu YC, Chang KT, Wu MT, Chiou YC, Su KC, Perng RP, Lee YC: **Leukotriene C4 induces TGF-beta1 production in airway epithelium via p38 kinase pathway.** *Am J Respir Cell Mol Biol* 2006, **34**:101–107.
63. Eap R, Jacques E, Semlali A, Plante S, Chakir J: **Cysteinyl leukotrienes regulate TGF-beta(1) and collagen production by bronchial fibroblasts obtained from asthmatic subjects.** *Prostaglandins Leukot Essent Fatty Acids* 2012, **86**:127–133.
64. Yamashita K, Gerken U, Vogel P, Hossmann K, Wiessner C: **Biphasic expression of TGF-beta1 mRNA in the rat brain following permanent occlusion of the middle cerebral artery.** *Brain Res* 1999, **836**:139–145.
65. Doyle KP, Cekanaviciute E, Mamer LE, Buckwalter MS: **TGFbeta signaling in the brain increases with aging and signals to astrocytes and innate immune cells in the weeks after stroke.** *J Neuroinflammation* 2010, **7**:62.

66. Kohta M, Kohmura E, Yamashita T: **Inhibition of TGF-beta1 promotes functional recovery after spinal cord injury.** *Neurosci Res* 2009, **65**:393–401.
67. Massague J, Wotton D: **Transcriptional control by the TGF-beta/Smad signaling system.** *EMBO J* 2000, **19**:1745–1754.
68. Kloos DU, Choi C, Wingender E: **The TGF-beta–Smad network: introducing bioinformatic tools.** *Trends Genet* 2002, **18**:96–103.
69. Wang H, Yang GH, Bu H, Zhou Q, Guo LX, Wang SL, Ye L: **Systematic analysis of the TGF-beta/Smad signalling pathway in the rhabdomyosarcoma cell line RD.** *Int J Exp Pathol* 2003, **84**:153–163.
70. ten Dijke P, Hill CS: **New insights into TGF-beta–Smad signalling.** *Trends Biochem Sci* 2004, **29**:265–273.
71. Levy L, Hill CS: **Smad4 dependency defines two classes of transforming growth factor beta (TGF-beta) target genes and distinguishes TGF-beta-induced epithelial-mesenchymal transition from its antiproliferative and migratory responses.** *Mol Cell Biol* 2005, **25**:8108–8125.
72. Kaminska B, Wesolowska A, Danilkiewicz M: **TGF beta signalling and its role in tumour pathogenesis.** *Acta Biochim Pol* 2005, **52**:329–337.
73. Finnson KW, Parker WL, Chi Y, Hoemann CD, Goldring MB, Antoniou J, Philip A: **Endoglin differentially regulates TGF-beta-induced Smad2/3 and Smad1/5 signalling and its expression correlates with extracellular matrix production and cellular differentiation state in human chondrocytes.** *Osteoarthr Cartil* 2010, **18**:1518–1527.
74. Chen M, Lv Z, Jiang S: **The effects of triptolide on airway remodelling and transforming growth factor-beta/Smad signalling pathway in ovalbumin-sensitized mice.** *Immunology* 2011, **132**:376–384.
75. Lampropoulos P, Zizi-Sermpetzoglou A, Rizos S, Kostakis A, Nikiteas N, Papavassiliou AG: **TGF-beta signalling in colon carcinogenesis.** *Cancer Lett* 2012, **314**:1–7.
76. Park BJ, Park JJ, Byun DS, Park JH, Chi SG: **Mitogenic conversion of transforming growth factor-beta1 effect by oncogenic Ha-Ras-induced activation of the mitogen-activated protein kinase signaling pathway in human prostate cancer.** *Cancer Res* 2000, **60**:3031–3038.
77. Dai C, Yang J, Liu Y: **Transforming growth factor-beta1 potentiates renal tubular epithelial cell death by a mechanism independent of Smad signaling.** *J Biol Chem* 2003, **278**:12537–12545.
78. Kim YK: **TGF-beta1 induction of p21WAF1/cip1 requires Smad-independent protein kinase C signaling pathway.** *Arch Pharm Res* 2007, **30**:739–742.
79. Niculescu-Duvaz I, Phanish MK, Colville-Nash P, Dockrell ME: **The TGFbeta1-induced fibronectin in human renal proximal tubular epithelial cells is p38 MAP kinase dependent and Smad independent.** *Nephron Exp Nephrol* 2007, **105**:e108–e116.
80. Kane NM, Jones M, Brosens JJ, Kelly RW, Saunders PT, Critchley HO: **TGFbeta1 attenuates expression of prolactin and IGFBP-1 in decidualized endometrial stromal cells by both SMAD-dependent and SMAD-independent pathways.** *PLoS One* 2010, **5**:e12970.
81. Watkins SJ, Borthwick GM, Oakenfull R, Robson A, Arthur HM: **Angiotensin II-induced cardiomyocyte hypertrophy *in vitro* is TAK1-dependent and Smad2/3-independent.** *Hypertens Res* 2012, **35**:393–398.
82. Gomes FC, Sousa Vde O, Romao L: **Emerging roles for TGF-beta1 in nervous system development.** *Int J Dev Neurosci* 2005, **23**:413–424.
83. Buss A, Pech K, Kakulas BA, Martin D, Schoenen J, Noth J, Brook GA: **TGF-beta1 and TGF-beta2 expression after traumatic human spinal cord injury.** *Spinal Cord* 2008, **46**:364–371.
84. Komuta Y, Teng X, Yanagisawa H, Sango K, Kawamura K, Kawano H: **Expression of transforming growth factor-beta receptors in meningeal fibroblasts of the injured mouse brain.** *Cell Mol Neurobiol* 2010, **30**:101–111.
85. Zhou Y, Fang SH, Ye YL, Chu LS, Zhang WP, Wang ML, Wei EQ: **Caffeic acid ameliorates early and delayed brain injuries after focal cerebral ischemia in rats.** *Acta Pharmacol Sin* 2006, **27**:1103–1110.
86. Zhang L, Zhang WP, Chen KD, Qian XD, Fang SH, Wei EQ: **Caffeic acid attenuates neuronal damage, astrogliosis and glial scar formation in mouse brain with cryoinjury.** *Life Sci* 2007, **80**:530–537.

doi:10.1186/1742-2094-9-145

Cite this article as: Huang et al.: Transforming growth factor beta1-induced astrocyte migration is mediated in part by activating 5-lipoxygenase and cysteinyl leukotriene receptor 1. *Journal of Neuroinflammation* 2012 **9**:145.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

